

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.



ISSN:2155-9899

Journal of Clinical & Cellular Immunology

The International Open Access

Journal of Clinical & Cellular Immunology

Special Issue Title: Cancer Immunology

Handling Editor

Dr. Haval Shirwan

University of Louisville, USA

Available online at: OMICS Publishing Group (www.omicsonline.org)

This article was originally published in a journal published by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/2155-9899.S5-004>

Exploring the Therapeutic Efficacy of Glioma Vaccines Based on Allo- and Syngeneic Antigens and Distinct Immunological Costimulation Activators

Apostolos Stathopoulos^{1,2,7*}, Chrystel Pretto², Laurent Devillers², Denis Pierre², Florence M. Hofman³, Alan L. Epstein³, Hooman Farghadani⁸, Carol A. Kruse⁴, Martin R. Jadus⁵, Thomas C. Chen^{2,7} and Virgil E.J.C. Schijns^{2,6*}

¹Department of Neurosurgery, Arlon Hospital, Arlon, Belgium

²Epitopeic Research Corporation (ERC), Namur, Belgium

³Department of Pathology, University of Southern California, Keck School of Medicine, Los Angeles, California, USA

⁴Department of Neurosurgery, University of California, Los Angeles, California and the Jonsson Comprehensive Cancer Center, UCLA, Los Angeles, CA 90095, USA

⁵Veterans Affairs Medical Center, Long Beach, CA 90822, box 113, 5901 E7th St. and Chao Cancer Center, University of California, Irvine, Orange CA, USA

⁶Cell Biology & Immunology Group, Wageningen University, PO Box 338, 6700 AH Wageningen, The Netherlands

⁷Department of Neurosurgery, University of Southern California, Keck School of Medicine, Los Angeles, California, USA

⁸Department of General Surgery, Arlon Hospital, Arlon, Belgium

Abstract

The efficacy of a various immunotherapeutic immunisation strategies for malignant glioma brain cancer was evaluated in the syngeneic CNS-1 Lewis rat glioma model. A prototype glioma cancer vaccine, which was composed of multivalent antigens derived from allogeneic and syngeneic cells and lysates, formed the prototype preparation of antigens. These antigens reflect the autologous antigens derived from the patient's surgically removed tumor tissue, as well as allogeneic antigens from glioma tumor tissue surgically removed from donor patients. This antigen mixture provides a broad spectrum of tumor associated antigens (TAA) and helps to prevent escape of tumor immune surveillance when given as a vaccine. This antigen preparation was administered in a therapeutic setting with distinct single or multiple co-stimulation-favouring immunostimulants and evaluated for inhibition of tumor growth. Our prototype vaccine was able to arrest progression of tumor growth when co-delivered in a specific regimen together with the costimulating multi-TLR agonist, Bacille Calmette Guerin (BCG) and interleukin-2, or with the Toll-Like receptor (TLR) 7/8 activator resiquimod.

Introduction

Glioblastoma multiforme (GBM) is an invasive malignant tumor of the central nervous system. Conventional therapy options include surgery, radiation, and chemotherapy, but with them the prognosis for GBM patients is limited to a mean survival time of only 14.6 months [1]. Immunotherapy is emerging as a novel complementary treatment option for a variety of malignancies including GBM. The use of successful passive immunotherapies based on the administration of immune elements, such as antibodies has proven very successful against various types of cancer. Well-known examples include antibodies that target tumor expressing receptors for epidermal growth factor (EGF), called Herceptin, those that target HER-2 [2], and those that are directed against angiogenic, tumor blood vessel growth-promoting vascular endothelial growth factor (VEGF), known as Avastin [3]. Recently, the antibody directed against an immune response inhibitory molecule, called cytotoxic T lymphocyte associated protein 4 (CTLA4), known as Yervoy[®], has shown promising clinical efficacy in melanoma patients [4].

Apart from these passive antibody-based therapies, a range of active immunotherapies are in late stage development and are close to reaching approval as standard of care. These clinical studies clearly demonstrate that the immune system is able to discriminate cancer cells from normal cells following recognition of tumor associated antigens (TAA). Indeed a recent approval for a prostate cancer vaccine named Provenge[®], was obtained from the FDA in April 2010, for the treatment of asymptomatic or minimally symptomatic metastatic, castrate-resistant (hormone-refractory) prostate cancer [5]. This recent approval has rejuvenated interest in the field as a whole.

In the present study we demonstrate a prototype brain cancer vaccine against gliomas which is composed of multivalent antigens derived from allogeneic and syngeneic cells and lysates. Our prototype is reflected in a clinical situation by autologous antigens derived from

the patient's surgically removed tumor tissue. In addition, glioma tumor tissue surgically removed from donor patients provides a second source of allogeneic antigens that can be isolated and subsequently stored for later use. This material provides a new source of TAA that may display HLA-restriction that may overlap with that on the patient's glioma. They may serve to enhance the overall immune response. If processed under Good Manufacturing Conditions, it may provide an "off-the-shelf" application. Relevant unique or shared TAAs overexpressed by tumor cells are present among thousands of irrelevant immunotolerant non-tumor associated antigens. The broad range of TAAs is preferred over vaccines with a mono- or oligo-valent antigenic content. These vaccines will prevent escape of tumor cells due to antigenic loss, or active MHC downregulation. In addition, a tumor antigen mixture also circumvents the use of monovalent synthetic peptides, which are restricted for use in patients with defined HLA types. By including TAAs of allogeneic origin, we additionally trigger allogeneic immune reactions. The haplotypes of CNS-1 and RG2 are fairly close, i.e., RT1^l vs RT1^{bl}, yet provide a mismatch that may be considered aberrant self,

***Corresponding authors:** Dr. Virgil E.J.C. Schijns, Cell Biology & Immunology Group, Wageningen University, PO Box 338, 6700 AH Wageningen, The Netherlands, E-mail: Virgil.schijns@wur.nl

Dr. Apostolos Stathopoulos, Department of Neurosurgery, Arlon Hospital, Arlon, Belgium, E-mail: tstath@hotmail.com

Received December 16, 2011; **Accepted** March 06, 2012; **Published** March 09, 2012

Citation: Stathopoulos A, Pretto C, Devillers L, Pierre D, Hofman FM, et al. (2012) Exploring the Therapeutic Efficacy of Glioma Vaccines Based on Allo- and Syngeneic Antigens and Distinct Immunological Costimulation Activators. J Clin Cell Immunol S5:004. doi:[10.4172/2155-9899.S5-004](http://dx.doi.org/10.4172/2155-9899.S5-004)

Copyright: © 2012 Stathopoulos A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and also induce an alloresponse. This may provide better protection due to the “non-self” immune recognition of these antigens.

Although allogeneic tumor antigens may provoke immune responses to non-self antigens in classical allogeneic immune rejections, glioma tumors, once established, are known to actively suppress the host’s immune system, by well characterized mechanisms [6], which often leads to subsequent evasion of immune surveillance. We therefore decided to test various signaling and costimulation-favouring immunostimulants in combination with our prototype vaccine antigen formulation for anti-tumor activity in an aggressive rat glioma brain tumor model, CNS-1, that is syngeneic in Lewis rats.

Here we demonstrate that our prototype vaccine is able to arrest progression of tumor growth when co-delivered in a specific regimen with the costimulatory-enhancing multi-TLR agonist, Bacille Calmette Guerin (BCG) adjuvant with interleukin-2, or with the TLR 7/8 activator resiquimod.

Materials and Methods

Tumor model

Glioma cells: Both CNS-1 tumor cells and RG2 (ATCC, CRL-2433) rat glioma cells were cultured in Dulbecco’s modified Eagle (DMEM) medium containing 10% bovine fetal serum (FBS), 5% antibiotics Penicillin, Streptomycin, and amphotericin B (Hyclone) in 175-mm² flasks. The CNS-1 haplotype is RT1^l and RG2 is RT1^{lv}.

Animals: Rat CNS-1 cells (2×10^5 cells/200 ul) were implanted subcutaneously (SC) using a 21 gauge needle into the right flank of 8-12 week-old (300 gram body weight) male Lewis rats. For each treatment group and control, 4-8 rats/group were used. All animal studies were approved by an independent ethical committee.

Monitoring tumor growth

The sizes of the CNS-1 tumor volumes were measured using a caliper three times per week on Mondays, Wednesdays and Fridays to monitor the effects of each treatment group.

Completion of experiment

Tumor implanted rats were sacrificed if they showed signs of discomfort, as defined by the ethical committee. For example if they appeared moribund due to weight loss, lethargy, ruffled fur, or when tumors showed ulceration. A mixture of Rompun and ketamine was used for anesthesia, followed by a dose of sodium pentobarbital for euthanasia.

Vaccine

The vaccine antigen preparation was composed of a mixture of haptized CNS-1 cells and RG2 cells (1×10^6 CNS-1 syngeneic and 1×10^6 RG2 allogeneic glioma cells), together with lysates produced from 3×10^6 CNS-1 syngeneic cells and 3×10^6 allogeneic RG2 glioma cells. The haptization method has been described before [7].

This vaccine preparation was kept as a constant factor and given in combination with various other anti-tumor or immunostimulatory agents as specified in Figures 1A & 2A. The TLR7/8 agonist resiquimod was co-administered 3 times a week on Mondays, Wednesdays and Fridays.

Chemicals and Reagents

Immunomodulators and potentiators

Rats were subcutaneously (SC) injected in the flank, contralateral

to the tumor-implanted side, with resiquimod (R848) (Invitrogen, TLRL-R848), a Toll-like receptor 7/8 agonist, at a dose of 100 µg/kg, corresponding to 30 µg/dose, three times per week on Mondays, Wednesdays and Fridays.

In a parallel arm of the experiment we evaluated the effect of cyclophosphamide (CY) administration, in diverse regimens and dosing as specified, on CNS-1 glioma development. Cyclophosphamide (CalBiochem, 239785) was given at 30 mg/kg 3 times per week, or at 100 mg/kg once every week (See also Figure 1A and 2A).

We also tested the vaccine antigen preparations combined with Bacilles Calmette-Guérin, interleukin-2 (IL-2), or a B7.1 fusion protein.

The Bacilles Calmette-Guérin (BCG), substrain Connaught, (Immunocyst[®], Sanofi Pasteur) is an agonist of TLRs 2, -4 and -9, was used at 2×10^5 CFU per dose.

Interleukin-2 (IL-2) was injected daily at a dose of 75,000 IU/day from Monday to Friday starting on the day of vaccine injection.

A B7.1 fusion protein consisting of the extracellular domains of human B7.1 and the Fc portion of human IgG1, called B7.1-Fc, was produced as described [8]. This protein induced complete regression of Colon 26 tumors in a mouse model and slowed tumor growth dramatically in mice with established poorly immunogenic RENCA and Madison109 tumors [8]. The B7.1-Fc protein was diluted in sterile PBS at 250 µg/200µl concentration.

For statistical analysis, we used ANOVA nonparametric testing followed by student’s *t* tests to compare groups. P values of < 0.05 (*), *p* < 0.01 (**), *p* < 0.001 (***) were considered statistically significant.







Figure 1: A vaccine was prepared from allogeneic and syngeneic antigens and administered in a fractionated regimen together with BCG and IL-2 to determine the effects on CNS-1 glioma growth. A) Treatment schedule representing one cycle of the various immunotherapeutic injections. B) Average values of tumor growth as a function of time post-implantation for the different treatment groups.

Results

Efficacy of a BCG-containing vaccine administered at day 17 after implantation

CNS-1 tumor cells (2×10^5 cells / 200 μ l) were implanted in syngeneic male Lewis rats and after 17 days were either left untreated (control group; n = 4) or administered with a therapeutic immunomodulatory regime or some of its components, as specified in the treatment schedule of Figure 1A.

The vaccine antigen preparation consisted of a mixture of haptized 10^6 syngeneic and 10^6 allogenic RG2 glioma cells together with lysates produced from 3×10^6 syngeneic CNS-1 cells and 3×10^6 allogenic RG2 gliomacells (n = 4). When monitoring tumor growth over time with a caliper, we observed no significant difference in tumor volumes between the control group (G1) and the rats receiving cyclophosphamide (CY) only at a dose of 100 mg/kg (G2) (Figure 1). From other experiments we had learned that the vaccine antigen preparation showed no anti-tumor efficacy by itself under the conditions described (for illustration see also figure 3, the antigen only group indicated by closed circles). By contrast, when this vaccine antigen preparation was administered together with the multiple TLR agonist BCG plus IL-2, an inhibition of tumor growth was noted (G6). Similarly, the same treatment schedule supplemented with CY (30 mg/kg on day 19, and 80 mg/kg on day 31) also inhibited tumor growth (G4). Remarkably, also five daily injections of B7.1-FC fusion



protein alone (Monday to Friday, starting day 17 post implantation), showed some inhibition of tumor growth (G3), though not statistically significant relative to control groups. However, when combining the beneficial vaccine preparation plus CY (30 mg/kg) treatment with an additional regimen of B7.1-Fc antibody injections no inhibition of tumor growth was noted (G5).

A BCG-containing vaccine shows better efficacy amongst the diverse vaccine prototypes tested

In a subsequent experiment, CNS-1 tumor cells (2×10^5 cells / 200 μ l) were implanted in 8-12 week-old syngeneic male Lewis rats and were either left untreated (control group; n = 4) or administered with a therapeutic immunomodulatory treatment regimen or some of its components, as specified in the treatment schedule depicted in Figure 2A. Instead of waiting until day 17 post tumor implantation, we now started at day 10 after tumor inoculation.

As before, the vaccine antigen preparation consisted of a mixture of haptized syngeneic CNS-1 and allogenic RG2 glioma cells (10^6 each) with lysates produced from syngeneic CNS-1 cells and allogenic RG2 glioma cells (3×10^6 each) (n = 4). When monitoring tumor growth over time with a caliper, we observed no significant difference in tumor volumes between the untreated control group (G1) and the rats receiving low dose cyclophosphamide (CY, 30 μ g/dose) in week two (G4), B7.1-Fc fusion protein only (G3), B7.1-Fc protein

plus CY (G2) or vaccine plus CY only (G5) (Figure 2A). By contrast, when this vaccine antigen preparation was administered together with the multiple TLR agonist BCG plus IL-2 and CY (100 µg/dose), a significant inhibition ($p < 0.05$) of tumor growth was noted (G6). This vaccine preparation was administered in a fractionated schedule starting with allogeneic cells on the first day, followed by syngeneic cells the second day, and allogeneic and syngeneic lysates on the third and fourth days, respectively (G6). Remarkably, also *ex vivo* tumor tissue isolated from a syngeneic established CNS-1 tumor plus CY (G7) showed some inhibition of tumor growth (although not statistically significant) relative to control groups (Figure 2B).

Therapeutic treatment with a vaccine containing immunostimulatory TLR7/8 agonist shows marked inhibition of glioma tumor growth

In view of the beneficial effect of vaccination in the context of the multi-TLR agonist BGC, we decided to evaluate the effect of a TLR7/8 agonist in our CNS-1 tumor model. CNS-1 tumor cells (2×10^5 cells/200 µl) were implanted in 8-12 week-old syngeneic male Lewis rats and either left untreated (control (CTRL) group; $n = 8$) or treated at day 10 after implantation with a vaccine antigen preparation consisting of a mixture of haptenized syngeneic CNS-1 and allogeneic RG2 glioma cells (10^6 each) with lysates produced from syngeneic CNS-1 and allogeneic RG2 glioma cells (3×10^6 each) ($n = 8$), given alone or combined with the TLR7/8 agonist, resiquimod (R848). This vaccine preparation was given 3 times a week on Mondays, Wednesdays and Fridays. When monitoring tumor growth over time with a caliper, we observed no significant difference in tumor growth between control and vaccine antigen only treated rats (Figure 3). By contrast, when this antigen preparation was administered together with a TLR7/8 agonist (30 µg/dose) significant inhibition ($p, 0.001, ***$) of tumor growth, with complete tumor regression was noted (Figure 3). Remarkably, administration of TLR7/8 agonist alone in the same regimen also inhibited tumor ($p, 0.001, ***$) (Figure 3).

Inhibition of tumor progression with large established glioma tumors by a vaccine containing the TLR7/8 immunostimulant

As mentioned above, Lewis rats treated with a vaccine without immunostimulatory TLR7/8 agonist showed little inhibition of tumor growth relative to untreated control rats, while the TLR7/8 containing vaccine strongly suppressed tumor growth when given 3 times per week starting at day 10 after implantation. We therefore decided to examine the effect of therapeutic vaccination of animals which showed no inhibition of tumor growth at day 34 after implantation. We administered the TLR7/8 containing vaccine to animals with large (12-14,000 mm³) tumors, starting at 38 days after tumor implantation. The TLR7/8 agonist-containing vaccine was injected 3 times a week on Mondays, Wednesdays and Fridays, in both untreated control rats ($n=4$) and rats immunized with identical antigen only ($n=4$). We noted in both groups a slower rate of growth of gliomas with large tumor volumes, but the TLR7/8 containing vaccine was best able to evoke a significant ($p < 0.001, ***$), arrest of tumor volume growth in the rats that were treated before with the antigen only vaccine preparation, relative to rats which were untreated (Figure 4). As well, no signs of vaccination-induced adverse effects or toxicity were observed, confirming data from another study that immunotherapy is well tolerated with limited toxicity [9]. These untoward effects point to the relative safety and tolerability of vaccines.

Discussion

In the present study we show that a prototype vaccine, consisting of a mixture of allogeneic and syngeneic glioma cells and their lysates, administered together with either a multiple TLR2, -4 and -9 agonist BCG [9], or the TLR7/8 agonist resiquimod, is able to inhibit CNS-1 glioma brain tumor growth in syngeneic Lewis rats. For early stage tumors, a complete regression of tumor growth volume was noted, while for large established 38-day old tumors (14,000 mm³ volume), inhibition of tumor growth was noted for the TLR7/8 containing vaccine. These data confirm the well-known phenomenon that large glioma tumors are more difficult to control relative to smaller tumors. Nevertheless, if the latter is most relevant to the large tumor burden that may be present in individuals with gliomas in unresectable sites (e.g. parietal or temporo-parietal tumors), instituting vaccination with TLR agonists may be valuable in providing additional quality survival time.

During glioma tumor development a proportional increase in local and systemic immunosuppression has been documented [6,9,11,12]. Gliomas are known to contain T regulatory cells, which normally act as suppressor cells, regulating homeostasis and preventing autoimmune reactions. However, regulatory T cells also inhibit T effector functions, activated naturally by vaccination [13,14], and thereby facilitating tumor immune evasion and subsequent tumor progression. At the molecular level this is associated with an increase in production of immune inhibitory cytokines including transforming growth factor (TGF)-beta, and/or interleukin-10 [15]. In general, immunosuppression can be counteracted by either strong stimulation of proinflammatory immune pathways, by blocking immune inhibitory elements triggered by the tumor itself, or by a combined "push-pull" approach [16]. We therefore evaluated well-known costimulation-enhancing immunostimulants, such as TLR-agonists [17], and inhibitors of immune system signals, such as what the B7.1 and cyclophosphamide immunomodulators provide. They were administered in conjunction with our standard vaccine antigen preparation.

We noted beneficial antitumor activity with daily injection of the B7.1Fc fusion protein starting at day 17 after tumor implantation as a monotherapy. This soluble costimulator protein was fused to the Fc portion of the antibody. It has been developed and tested for the immunotherapy of solid tumors in mouse tumor models [8]. This B7.1-Fc protein was found to be capable of activating T cells *in vitro* and *in vivo* [8]. B7.1 is able to engage with two known receptors. CD28, which triggers a stimulatory signal to activate naive T cells after binding by B7.1 (4), and its counterreceptor, CTLA-4, which triggers a negative signal to stop T-cell activation. Since B7.1 has high affinity to CTLA-4, soluble B7.1-Fc may block CTLA-4 signaling instead of cross-linking CTLA-4, thereby sustaining the activation of tumor-specific T cells [18,19]. Crosslinking of T cell activating CD28 may be less important than blocking the inhibitory CTLA-4, since activated T cells recognizing tumor antigens requires less costimulation [20]. When B7.1 antibody was combined with other immune system activating treatments, we observed no synergistic activities, which illustrates that the timing and choice of modalities during combined immunotherapy is critical for a beneficial antitumor effect.

Also, cyclophosphamide, when dosed carefully, has been demonstrated to facilitate immunotherapy of established tumors through the elimination/inactivation of suppressor T cells [21-23]. In our investigations, a protocol of cyclophosphamide (CY)

only, administered at a dose of 100 mg/kg given on day 19 post transplantation did not affect tumor growth (Figure 1), while three lower doses of 30 mg/kg each starting on day 17 given within a week period, did not show a beneficial anti-tumor activity (Figure 2). By contrast, CY treatment alone may even exacerbate tumor growth, relative to the untreated control group, which may likely result from suppression of endogenous anti-tumor immune reactions in parallel to inadequate direct tumor growth inhibition, resulting in a net increase in tumor growth volume. Nevertheless, when combined with other immune adjuvants or vaccines, beneficial effects were noted despite co-administration of CY.

The best efficacy was noted in treatment schedules containing the vaccine antigen preparation plus BCG and IL-2. *Bacillus Calmette-Guérin* (BCG) is a mycobacterium which contains various agonists of Toll-like receptor 2 (TLR2) and TLR4 in its cell wall skeleton (CWS), and also BCG DNA which acts as a TLR9 agonist [10,24]. Intravesical immunotherapy with BCG, as monotherapy, causes significant reductions in cancer tumor progression and is currently the standard treatment for superficial bladder cancer [25]. However, the exact mode of action of successful BCG therapy remains elusive, although massive cytokine expression and influx of innate immune cells have been documented, which may be involved in tumor elimination.

Apart from BCG, we also examined the effect of the TLR7/8 agonist resiquimod as an adjuvant in a separate animal study. TLR7/8 agonists exert pleiotropic effects on various immune cells which leads to stimulation of proinflammatory cytokine and chemokine production, as well as up-regulation of costimulatory molecules on antigen presenting cells [26]. The TLR7/8 agonist resiquimod has been shown to act as a vaccine adjuvant in investigational vaccine models [25,27], and to generate clinical-grade mature DCs [28].

In our CNS-1 model, the TLR7/8 agonist resiquimod showed superior antitumor effects against early tumor growth and remarkably, administration of TLR7/8 agonist alone inhibited tumor growth. The exact mode of action for the observed resiquimod-mediated anti-tumor immunity needs to be defined in detailed follow-up studies. Most likely the observed resiquimod-based immunotherapy, even in absence of additional tumor antigens, is able to activate a spontaneous, natural, innate anti-tumor immune response, that under normal circumstances is unable to control tumor growth. Hence, such dedicated follow-up studies need to address to involvement of anti-tumor killer macrophages or NK cells, or IFNs for the resiquimod-induced glioma growth regression. In addition, the TLR7/8 vaccine preparation evoked an arrest of tumor volume growth in rats carrying large tumors that were treated before with the antigen only vaccine preparation.

Although the presented results are promising the should always be interpreted with caution. Subcutaneous tumors may be exposed to markedly distinct immune conditions than in the brain, and so the physiologic as well as clinical relevance has to be validated in further studies.

Our vaccine preparation contains multiple tumor associated antigens (TAA) prepared from syngeneic and allogeneic cells. Although the precise identity of antigens is unknown we chose this antigen preparation for our vaccination strategy, rather than one or a few molecular targets, in order to target multiple TAAs simultaneously. This approach minimizes the chances for classical tumor-escape of immune control, as a result of selective growth of antigen-loss variants. With this approach we guaranteed that multiple relevant TAAs gain

enhanced exposure to the immune system, while the self-antigens in this mixture are neglected as a result of immunological tolerance. By including cell associated antigens, we increased the chance of cross-presentation by MHC class I molecules for priming of cytolytic CD8⁺ T cells. By including TAAs of allogeneic origin, we additionally trigger allogeneic immune reactions. The haplotypes of CNS-1 and RG2 are fairly close, i.e., RT1^l vs RT1^h thus providing a mismatch that may be considered aberrant self, and also induce a vigorous alloresponse. Allogeneic (major histocompatibility complex MHC-mismatched) tumor cells have shown better protection in a number of models likely due to the “non-self” immune recognition of these antigens [29-31].

Immunotherapy against TAA theoretically carries the risk of autoimmune reactions. Autoimmunity is possible, especially in view of our manipulations aimed at blocking immune response inhibitors coupled with the use of immunostimulating adjuvants. In the present studies, however, we have not observed any adverse reactions using our animal well-being scoring system. These observations are in line with well-documented observations, that therapeutic tumor vaccination is, in general, well-tolerized with minimal or no adverse events.

In conclusion, we noted beneficial antitumor activity of daily injection of the B7.1-Fc fusion protein as monotherapy, and with vaccine preparations that were administered with BCG and IL-2 only, or with the TLR7/8 agonist resiquimod. Altogether these data illustrate the importance of a beneficial immunomodulatory protocol. Different immunomodulators may result in distinct antitumor efficacy. Hence, we conclude from our data that only specific combinations of the right antigenic mixtures in conjunction with a suitable immunopotentiator is able to arrest aggressive glioma tumor growth in this experimental tumor model. Our data highlight the need for continued, more extensive exploration into such combinations.

References

1. Stupp R, Roila F, ESMO Guidelines Working Group (2009) Malignant glioma: ESMO clinical recommendations for diagnosis, treatment and follow-up. *Ann Oncol* 4: 126-128.
2. Hudis CA (2007) Trastuzumab-mechanism of action and use in clinical practice. *N Engl J Med* 357: 39-51.
3. Ananthnarayan S, Bahng J, Roring J, Nghiemphu P, Lai A, et al. (2008) Time course of imaging changes of GBM during extended bevacizumab treatment. *J Neurooncol* 88: 339-347.
4. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, et al. (2010) Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 363: 711-723.
5. Buonerba C, Ferro M, Di Lorenzo G (2011) Sipuleucel-T for prostate cancer: the immunotherapy era has commenced. *Expert Rev Anticancer Ther* 11: 25-28.
6. Gomez GG, Kruse CA (2006) Mechanisms of malignant glioma immune resistance and sources of immunosuppression. *Gene Ther Mol Biol* 10: 133-146.
7. Henry N, Claman HN, Miller SD (1976) Requirements for Induction of T Cell Tolerance to Dnfb: Efficiency of Membrane-Associated DNFB. *J Immunol* 117: 480-485.
8. Liu A, Hu P, Khawli LA, Epstein AL (2005) Combination B7-Fc Fusion Protein Treatment and Treg Cell Depletion Therapy. *Clin Cancer Res* 11: 8492-8502.
9. de Vleeschouwer S, Rapp M, Sorg RV, Steiger HJ, Stummer W, et al. (2006) Dendritic cell vaccination in patients with malignant gliomas: current status and future directions. *Neurosurgery* 59: 988-999.
10. Uehori J, Matsumoto M, Tsuji S, Akazawa T, Takeuchi O, et al. (2003) Simultaneous blocking of human Toll-like receptors 2 and 4 suppresses

- myeloid dendritic cell activation induced by *Mycobacterium bovis* bacillus Calmette-Guérin peptidoglycan. *Infect Immun* 71: 4238-4249.
11. Dix AR, Brooks WH, Roszman TL, Morford LA (1999) Immune defects observed in patients with primary malignant brain tumors. *J Neuroimmunol* 100: 216-32.
 12. Ge L, Hoa N, Bota DA, Natividad J, Howat A, et al. (2010) Immunotherapy of brain cancers: the past, the present, and future directions. *Clin Dev Immunol* 2010: 296453.
 13. El Andaloussi A, Lesniak MS (2006) An increase in CD4+CD25+FOXP3+ regulatory T cells in tumor-infiltrating lymphocytes of human glioblastoma multiforme. *Neuro Oncol* 8: 234-243.
 14. Fecci PE, Mitchell DA, Whitesides JF, Xie W, Friedman AH, et al. (2006) Increased regulatory T-cell fraction amidst a diminished CD4 compartment explains cellular immune defects in patients with malignant glioma. *Cancer Res* 66: 3294-3302.
 15. Qiu B, Zhang D, Wang C, Tao J, Tie X, et al. (2011) IL-10 and TGF- β 2 are overexpressed in tumor spheres cultured from human gliomas. *Mol Biol Rep* 38: 3585-3591.
 16. Ahlers JD, Belyakov IM, Terabe M, Koka R, Donaldson DD, et al. (2002) A push-pull approach to maximize vaccine efficacy: abrogating suppression with an IL-13 inhibitor while augmenting help with granulocyte/macrophage colony-stimulating factor and CD40L. *Proc Natl Acad Sci U S A* 99: 13020-13025.
 17. Kawai T, Akira S (2007) TLR signaling. *Semin Immunol* 19: 24-32.
 18. Sturmhoefel K, Lee K, Gray GS, Thomas J, Zollner R, et al. (1999) Potent activity of soluble B7-IgG fusion proteins in therapy of established tumors and as vaccine adjuvant. *Cancer Res* 59: 4964 - 4972.
 19. Moro M, Gasparri AM, Pagano S, Bellone M, Tornaghi P, et al. (1999) Induction of therapeutic T-cell immunity by tumor targeting with soluble recombinant B7-immunoglobulin costimulatory molecules. *Cancer Res* 59: 2650-2656.
 20. Croft M, Bradley LM, Swain SL (1994) Naive versus memory CD4 T cell response to antigen. Memory cells are less dependent on accessory cell costimulation and can respond to many antigen-presenting cell types including resting B cells. *J Immunol* 152: 2675-2685.
 21. North RJ (1982) Cyclophosphamide-facilitated adoptive immunotherapy of an established tumor depends on elimination of tumor-induced suppressor T cells. *J Exp Med* 155: 1063-1074.
 22. Ghiringhelli F, Larmonier N, Schmitt E, Parcellier A, Cathelin D, et al. (2004) CD4+CD25+ regulatory T cells suppress tumor immunity but are sensitive to cyclophosphamide which allows immunotherapy of established tumors to be curative. *Eur J Immunol* 34: 336-344.
 23. Ghiringhelli F, Menard C, Puig PE, Ladoire S, Roux S, et al. (2007) Metronomic cyclophosphamide regimen selectively depletes CD4+CD25+ regulatory T cells and restores T and NK effector functions in end stage cancer patients. *Cancer Immunol Immunother* 56: 641-648.
 24. Hoppstädter J, Diesel B, Zarbock R, Breinig T, Monz D, et al. (2010) Differential cell reaction upon Toll-like receptor 4 and 9 activation in human alveolar and lung interstitial macrophages. *Respir Res* 11: 124.
 25. Lamm DL (1992) Long-term results of intravesical therapy for superficial bladder cancer. *Urol Clin North Am* 19: 573-580.
 26. Smits EL, Ponsaerts P, Berneman ZN, Van Tendeloo VF (2008) The use of TLR7 and TLR8 ligands for the enhancement of cancer immunotherapy. *Oncologist* 13: 859-875.
 27. Tomai MA, Vasilakos JP (2011) TLR-7 and -8 agonists as vaccine adjuvants. *Expert Rev Vaccines* 10: 405-407.
 28. Boullart AC, Aarntzen EH, Verdijk P, Jacobs JF, Schuurhuis DH, et al. (2008) Maturation of monocyte-derived dendritic cells with Toll-like receptor 3 and 7/8 ligands combined with prostaglandin E2 results in high interleukin-12 production and cell migration. *Cancer Immunol Immunother* 57: 1589-1597.
 29. Knight BC, Souberbielle BE, Rizzardi GP, Ball SE, Dalgleish AG (1996) Allogeneic murine melanoma cell vaccine: a model for the development of human allogeneic cancer vaccine. *Melanoma Res* 6: 299-306.
 30. Hrouda D, Todryk SM, Perry MJ, Souberbielle BE, Kayaga J, et al. (2000) Allogeneic whole-tumour cell vaccination in the rat model of prostate cancer. *BJU Int* 86: 742-748.
 31. Spivey TL, Uccellini L, Ascierio ML, Zoppoli G, De Giorgi V, et al. (2011) Gene expression profiling in acute allograft rejection: challenging the immunologic constant of rejection hypothesis. *J Transl Med* 9: 174.

This article was originally published in a special issue, **Cancer Immunology** handled by Editor(s). Dr. Haval Shirwan, University of Louisville, USA

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 200 Open Access Journals
- 15,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.editorialmanager.com/clinicalgroup